

# Asthma Treatment Plan

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)

The Pediatric/Adult  
Asthma Coalition  
of New Jersey

"Your Pathway to Asthma Control"  
PACNJ approved Plan available at  
www.pacnj.org

Sponsored by  
**AMERICAN LUNG ASSOCIATION**  
IN NEW JERSEY



(Please Print)

Name		Date of Birth	Effective Date
Doctor	Parent/Guardian (if applicable)		Emergency Contact
Phone	Phone		Phone

## HEALTHY



You have all of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above \_\_\_\_\_

**Take daily medicine(s). Some metered dose inhalers may be more effective with a "spacer" – use if directed.**

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Advair® <input type="checkbox"/> 100, <input type="checkbox"/> 250, <input type="checkbox"/> 500	_____ 1 inhalation twice a day
<input type="checkbox"/> Advair® HFA <input type="checkbox"/> 45, <input type="checkbox"/> 115, <input type="checkbox"/> 230	_____ 2 puffs MDI twice a day
<input type="checkbox"/> Alvesco® <input type="checkbox"/> 80, <input type="checkbox"/> 160	_____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day
<input type="checkbox"/> Asmanex® Twisthaler® <input type="checkbox"/> 110, <input type="checkbox"/> 220	_____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations <input type="checkbox"/> once or <input type="checkbox"/> twice a day
<input type="checkbox"/> Flovent® <input type="checkbox"/> 44, <input type="checkbox"/> 110, <input type="checkbox"/> 220	_____ 2 puffs MDI twice a day
<input type="checkbox"/> Flovent® Diskus® <input type="checkbox"/> 50 <input type="checkbox"/> 100 <input type="checkbox"/> 250	_____ 1 inhalation twice a day
<input type="checkbox"/> Pulmicort Flexhaler® <input type="checkbox"/> 90, <input type="checkbox"/> 180	_____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations <input type="checkbox"/> once or <input type="checkbox"/> twice a day
<input type="checkbox"/> Pulmicort Respules® <input type="checkbox"/> 0.25, <input type="checkbox"/> 0.5, <input type="checkbox"/> 1.0	_____ 1 unit nebulized <input type="checkbox"/> once or <input type="checkbox"/> twice a day
<input type="checkbox"/> Qvar® <input type="checkbox"/> 40, <input type="checkbox"/> 80	_____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day
<input type="checkbox"/> Singulair <input type="checkbox"/> 4, <input type="checkbox"/> 5, <input type="checkbox"/> 10 mg	_____ 1 tablet daily
<input type="checkbox"/> Symbicort® <input type="checkbox"/> 80, <input type="checkbox"/> 160	_____ <input type="checkbox"/> 1, <input type="checkbox"/> 2 puffs MDI twice a day
<input type="checkbox"/> Other	
<input type="checkbox"/> None	

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take this medicine \_\_\_\_\_ minutes before exercise.

## Triggers

Check all items that trigger patient's asthma:

- Chalk dust
- Cigarette Smoke & second hand smoke
- Colds/Flu
- Dust mites, dust, stuffed animals, carpet
- Exercise
- Mold
- Ozone alert days
- Pests - rodents & cockroaches
- Pets - animal dander
- Plants, flowers, cut grass, pollen
- Strong odors, perfumes, cleaning products, scented products
- Sudden temperature change
- Wood Smoke
- Foods:

Other: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

## CAUTION



You have any of these:

- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other: \_\_\_\_\_

And/or Peak flow from \_\_\_\_\_ to \_\_\_\_\_

**Continue daily medicine(s) and add fast-acting medicine(s).**

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Accuneb® <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg	_____ 1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol <input type="checkbox"/> 1.25, <input type="checkbox"/> 2.5 mg	_____ 1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil®	_____ 2 puffs MDI every 4 hours as needed
<input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex®	_____ 2 puffs MDI every 4 hours as needed
<input type="checkbox"/> Xopenex® <input type="checkbox"/> 0.31, <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg	_____ 1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Increase the dose of, or add:	
<input type="checkbox"/> Other	

➔ If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.

## EMERGENCY



Your asthma is getting worse fast:

- Fast-acting medicine did not help within 15-20 minutes
- Breathing is hard and fast
- Nose opens wide
- Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue

And/or Peak flow below \_\_\_\_\_

**Take these medicines NOW and call 911. Asthma can be a life-threatening illness. Do not wait!**

<input type="checkbox"/> Accuneb® <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg	_____ 1 unit nebulized every 20 minutes
<input type="checkbox"/> Albuterol <input type="checkbox"/> 1.25, <input type="checkbox"/> 2.5 mg	_____ 1 unit nebulized every 20 minutes
<input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil®	_____ 2 puffs MDI every 20 minutes
<input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex®	_____ 2 puffs MDI every 20 minutes
<input type="checkbox"/> Xopenex® <input type="checkbox"/> 0.31, <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg	_____ 1 unit nebulized every 20 minutes
<input type="checkbox"/> Other	

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association of New Jersey, and PACNJ are supported by a grant from the New Jersey Department of Health and Senior Services (NJHSS), with funding provided by the U.S. Centers for Disease Control and Prevention (USCDC) under Cooperative Agreement #4U59CE000226-2. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NJHSS or the USCDC.

Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreement #487556017 to the American Lung Association of New Jersey, it does not constitute an EPA policy, nor does it represent the views of the Agency and no official endorsement should be inferred.

**REVISED MAY 2009**

Permission to reproduce blank form  
www.pacnj.org

### FOR MINORS ONLY:

- This student is capable and has been instructed in the proper method of self-administering of the non-nebulized inhaled medications named above in accordance with NJ Law.
- This student is not approved to self-medicate.

Make a copy for patient and for physician file. For children under 18, send original to school nurse or child care provider.

PHYSICIAN/APN/PA SIGNATURE \_\_\_\_\_ DATE \_\_\_\_\_

PARENT/GUARDIAN SIGNATURE \_\_\_\_\_

PHYSICIAN STAMP